

## CPD QUESTIONNAIRE 5.1.1

### Article: The management of recession midfacial to immediately placed implants in the aesthetic zone. Gluckman and Du Toit, page 6

1. *At immediate implant placement in the aesthetic zone:*
  - a Provisionalization or customizing of the abutment should not be done, the soft tissue needs first heal over the implant
  - b Provisionalization or customizing of the abutment should be done, the coronal tissues need supportin
2. *A narrow emergence profile allows for:*
  - a A greater soft tissue seal around the implant restoration/abutment
  - b Less displacement of the tissues coronal to the implant
  - c Greater soft tissue fill of the concavity around the implant restoration/abutment
  - d All the above
3. *Use of an internal, conical, Morse taper connection implant allows for:*
  - a Improved seal around the implant crown-abutment interface
  - b A reduction in micro-movement and microbial colonization of the implant-abutment interface
  - c Less displacement of the tissues coronal to the implant
  - d All the above
4. *The midfacial recession was salvageable in this case by soft tissue augmentation alone because:*
  - a The interproximal bony peaks and papillae at the implants were absent
  - b The interproximal bony peaks and papillae at the implants were augmented
  - c The interproximal bony peaks and papillae at the implants were intact
  - d The interproximal bony peaks and papillae at the implants had receded

5. *Following tooth extraction, loss of bone (especially on the buccofacial aspect as was seen in this case) is a result of:*
  - a Loss of the periodontal ligament that vascularizes bundle bone
  - b Loss of a coronal seal to the tooth socket
  - c Loss of the whole blood clot from which the alveolar socket heals
  - d Loss of the socket shield that supports the periodontal ligament and bundle bone

### Article: Making sense of mouth ulceration. Parts 5 and 6. Scully, page 24/28

6. *According to the author, main gastrointestinal disorders that may present with mouth ulceration include coeliac disease, and Crohn's and related diseases:*
  - a False
  - b True
7. *People with Coeliac disease have sensitivity to:*
  - a Casein
  - b Gelatin
  - c Gluten
8. *Which statement is correct. Patients suffering from Melkerrson-Rosenthal syndrome experience:*
  - a Lip swelling, Fissured tongue, Facial palsy
  - b Lip swelling alone
9. *Which is not a cause of lichenoid lesions?*
  - a Drugs, especially NSAIDs
  - b Diabetes
  - c Chemical burn
  - d Dental restorative materials
10. *Lichen Planus has a malignant potential of:*
  - a Over 3%
  - b Less than 1-3%
  - c More than 30%

## CPD QUESTIONNAIRE 5.1.2

### Article: Therapeutic Aesthetics. Greenwall, page 32

11. *What does carbamide peroxide break down into:*
  - a Urea and hydrogen
  - b Urea and water
  - c Urea and ammonia
12. *Which of the following materials is not expressly suggested for use in therapeutic aesthetics?*
  - a Carbamide peroxide
  - b Fluoride
  - c PRGF material
  - d Potassium nitrate
13. *What does the author state as an additional benefit of tooth whitening?*
  - a Reduced gingival inflammation
  - b Reduced plaque formation
  - c Improved gingival index scores
  - d All of the above
  - e None of the above
14. *Which patients can benefit from therapeutic aesthetics?*
  - a Elderly patients
  - b Dry mouth sufferers
  - c Chemotherapy and radiation patients
  - d All of the above
  - e None of the above
15. *Which statements are correct:*
  - a A concentration of 10% carbamide peroxide kills lactobacillus
  - b Chlorhexidene kills lactobacillus
  - c Chlorhexidine kills streptococcus mutans bacteria
  - d a and c
  - e b

### Article: Using injectable resin composite: parts one and two. Terry and Powers, page 52/64

16. *Some of the applications of the injectable resin composite technique include:*
  - a Paediatric composite crowns
  - b Transitional composite restorations (class III, IV, veneers)
  - c Establishing incisal edge length prior to aesthetic crown lengthening
  - d All of the above
  - e None of the above
17. *Which statement is correct:*
  - a The injectable resin composite technique is irreversible
  - b The injectable resin composite technique is reversible
18. *Which statement is incorrect:*
  - a The injectable resin composite technique can be performed without anaesthesia
  - b The injectable resin composite technique cannot be performed without anaesthesia
19. *According to the author, the major reasons for the restoration of primary teeth with full coronal coverage are:*
  - a Trauma
  - b Extensive caries
  - c Neither of the above
  - d Both of the above
20. *Treatment challenges associated with full coverage crowns in the past include:*
  - a Poor retention
  - b Gingival inflammation
  - c Fractured facings
  - d All of the above
  - e None of the above