

## CPD QUESTIONNAIRE 4.4.1

### Article: Periodontal and peri-implant tissue management in the esthetic zone. Verdecchia, page 14

- 1 Which statement is correct: In the case described, the patient presented for a vertical fracture of:
  - a The maxillary left lateral incisor
  - b The maxillary left central incisor
  - c The maxillary right central incisor
- 2 Which statement is correct: Horizontal and vertical bone defects were detected with a distance of:
  - a 4 mm from bone level to the contact points
  - b 5 mm from bone level to the contact points
  - c 6 mm from bone level to the contact points
- 3 Which statement is correct: To improve the stability of the membrane:
  - a A "double-layer" technique was used
  - b A "single-layer" technique was used
- 4 The period of extended interval allowed by the stability of the provisional bridge was:
  - a 3 months
  - b 6 months
  - c 4 months
- 5 Which statement is correct:
  - a A cement-retained provisional crown remained in situ 6 months
  - b A screw-retained provisional crown remained in situ 6 months
  - c A screw-retained provisional crown remained in situ 3 months
  - d A cement-retained provisional crown remained in situ 3 months

### Article: Making sense of mouth ulceration: part four. Scully, page 28

6. Which is not a predisposing factor in ANUG?
  - a Smoking
  - b Excessive alcohol consumption
  - c Immune defects, such as in HIV/AIDS
  - d. Viral respiratory infections
7. Syphilis is a chronic bacterial infection caused by:
  - a Mycobacterium tuberculosis
  - b Treponema pallidum
  - c Neither of the above
8. The incidence of TB is increasing annually by:
  - a 5%
  - b 8%
  - c 10%
9. Diagnosis of TB is by:
  - a Imaging
  - b Biopsy
  - c Microbial studies
  - d None of the above
  - e All of the above
10. To be considered cured of TB, a patient must have:
  - a Normal temperature
  - b Absence of ulceration
  - c. Three consecutive negative sputum cultures

## CPD QUESTIONNAIRE 4.4.2

### Article: Is alveolar ridge preservation (socket grafting) an effective therapy to preserve physiologic bone loss after tooth extraction?. Hartshorne, page 32

11. Which statement is incorrect: Eligibility criteria for the studies included:
  - a Studies that involved the application of any additional therapy that could have affected healing outcomes, e.g., simultaneous soft tissue grafting
  - b RCTs conducted in human subjects, older than 18 years, who had at least 1 nonmolar tooth extracted in which investigators compared ARP via socket filling
12. Which statement is correct:
  - a Absorbable collagen membrane was used in all the studies
  - b nonabsorbable polytetrafluoroethylene membrane was used in all the studies
13. Which statement is correct:
  - a In all 8 studies, the risk of bias due to inadequate random sequence generation and allocation concealment was considered to be high.
  - b In all 8 studies, the risk of bias due to inadequate random sequence generation and allocation concealment was considered to be low.
14. Which statement is correct: The limited number of studies used in the subgroup analysis found that:
  - a Flap elevation, use of a xenograft or an allograft, and the application of a membrane, had superior outcomes in terms of the horizontal dimensions only
  - b Flap elevation, use of a xenograft or an allograft, and the application of a membrane, had superior outcomes in terms of the vertical dimensions only
15. This review did not report on:
  - a Availability of bone volume
  - b Feasibility of implant placement at re-entry
  - c Whether further GBR or ridge expansion was required in the experimental and control groups at the implant placement stage
  - d None of the above
  - e All of the above

### Article: The management of immediate implant placement to optimize aesthetic outcome in the anterior maxilla. Gluckman and Du Toit, page 48

16. The appropriate flap for a CT graft at immediate implant placement without a socket defect is:
  - a Azzi's tunnel and pouch, partial-thickness flap
  - b Partial-thickness, with release incisions
  - c Full-thickness, with release incisions
  - d A semi-lunar flap
17. Steigman's recommendations for provisionalization propose:
  - a A feldspathic, true-to-life provisional
  - b A veneered composite, true-to-life provisional
  - c A polycarbonate, true-to-life provisional
  - d A natural tooth, true-to-life provisional
18. Covani recommends for immediate implant placement:
  - a The implant placed deeper, in the anatomical center of the extraction socket
  - b The implant placed deeper, lingual / palatal in the extraction socket
  - c the implant placed deeper, buccal / facial in the extraction socket
19. Selecting which implant may encourage bone growth toward the implant table:
  - a A morse taper internal connection
  - b An external hex connection
  - c An internal hex connection
  - d A tissue level single component
20. To compensate for buccofacial resorption, an immediate implant should be placed a minimum of:
  - a 4 mm from the socket's facial perimeter
  - b 3 mm from the socket's facial perimeter
  - c 2 mm from the socket's facial perimeter