

- 1 Which of the following statements is correct?
In the anterior tooth region, retention of or regeneration of a papillary height of 4 - 5 mm can be achieved if the inter-implant distance is at least:

a 2mm	b 3mm
c 4mm	d 5 mm
- 2 Which of the following statements is correct?:
Optimal papillary form is achieved when the distance between the bony crest and the connection point between an implant-borne crown and a pontic is

a 4.5 mm	b 3.4 mm
c 5.5 mm	d 6.3 mm
- 3 Which of the following statements is correct?
To obtain ideal implant aesthetics,

a the papilla height and form should be restored to correspond to that observed with natural teeth
b the papilla height and form should not correspond to that observed with natural teeth
c the papilla height but not the form should be restored to correspond to that observed with natural teeth
- 4 Which of the following statements is correct?

a The peri-implant gingiva, particularly if it is narrow, retracts 3 months after the abutment connection and restoration
b The peri-implant gingiva, particularly if it is narrow, retracts 6 months after the abutment connection and restoration
c The peri-implant gingiva, particularly if it is narrow, retracts 12 months after the abutment connection and restoration
d none of the above
- 5 Which of the following statements is correct?

a Only titanium abutments are recommended because hemidesmosomes have been shown to attach to them	
b Only zirconium abutments are recommended because hemidesmosomes have been shown to attach to them	
c Only gold abutments are recommended because hemidesmosomes have been shown to attach to them	
d a and b	e a and c
f b and c	g all of the above
h none of the above	
- 6 Which of the following statements is correct?

a It has not been proven that an excessive occlusal load during function can cause the loss of peri-implant bone
b It has been proven that an excessive occlusal load during function can cause the loss of peri-implant bone
c It has been proven that a decreased occlusal load during function can cause the loss of peri-implant bone
- 7 Which of the following statements is correct?:

a Soft-tissue recession around a wide - diameter implant averages 2.62 mm compared to 0.93 mm around a standard-diameter implant
b Soft-tissue recession around a wide - diameter implant averages 3.69 mm compared to 1.52 mm around a standard-diameter implant
c Soft-tissue recession around a wide - diameter implant averages 1.58 mm compared to 0.57 mm around a standard-diameter implant
d Soft-tissue recession around a wide - diameter implant averages 0.57 mm compared to 1.63 mm around a standard-diameter implant
- 8 Which of the following statements is correct?:
Optimal papillary form is achieved when the distance from the interdental bony crest to the contact point between the natural crown and the implant-borne is


a 4.5 mm	b 3.4 mm
c 5.5 mm	d 6.3 mm
- 9 Which of the following statements is correct?

a a thin biotype is generally more susceptible to peri-implant recession
b a thick biotype is generally more susceptible to peri-implant recession
c a thin biotype and a thick biotype are equally susceptible to peri-implant recession
- 10 Which of the following statements is correct?

a The biologic width is located subcrestally, and it is almost always subcrestal around the implant
b The biologic width is located supracrestally, and it is almost always supracrestal around the implant
c The biologic width is located supracrestally, and it is almost always subcrestal around the implant
d none of the above



MDL™
MINI DRIVE-LOCK



- IMMEDIATE LOAD
- STABILIZE DENTURES
- SIMPLIFIED PROCEDURES
- PROVISIONAL PROSTHETICS
- DIRECT PLACEMENT

To learn more about this product, please contact your local Implant Support Services sales consultant or customer services at:
Tel: 012 644 1040 , 021 851 4875
Fax: 012 644 1041
Email: bruce@implant.co.za

