## Article: BIO-ESTHETICS: giving a new face to smile enhancements. Dietschi, page 6

## 1. Which statement is correct:

- Improved dental esthetics and durable results have been for long intimately linked to ceramic restorations
- Improved dental esthetics and durable results have been for long intimately linked to modern composite resin technology

#### Which statement is correct: According to the author: 2.

- Resin composites are universally considered the "standard of care" material for the filling of small to medium class III, IV and V cavities
- Resin composites are universally considered the "standard of care" material for the filling of small to medium class I and II cavities

### 3. Which statement is correct: The new concept of 'bio-aesthetics' gives priority to additive, minimally or microinvasive procedures to preserve:

- Tooth biology а
- b Biomechanics
- Neither of the above C
- d Both of the above

#### 4. Which statement is incorrect:

- The majority of cases lead to evident solutions (direct and respectively indirect
- Conditions such as minor aesthetic form and color corrections or extensive decays in non-vital teeth, lead to evident solutions (direct and respectively indirect restorations)
- Neither of the above d Both of the above
- 5. Which statement is incorrect: According to the author, minimally invasive procedures include:
- Crowns and bridges
- Direct bonding, Ultra-thin Veneers, Modern inlays and onlay technique

## Article: Ceramics overview: classification by microstructure and processing methods. McLaren and Giordano, page 18

- Which statement is incorrect: According to the authors, ceramics are:
- Extremely brittle and will catastrophically fail after minor flexure. а
- A mixture of glass and crystal components.
- Strong in compression but weak in tension
- The flexural strength of Category 1: Glass-based systems, is usually in
- 160-170 MPa range
- 60-70 MPa range h
- 260-270 MPa range
- Which statement is correct: Zirconia has a fracture toughness reported to lie between:
- The magnitude of the shrinkage stresses generated from polymerisation shrinkage is less for most anterior composite restorations
- The magnitude of the shrinkage stresses generated from polymerisation shrinkage is less for most posterior composite restorations

#### 9. Which statement is correct:

- Alumina-based ceramics are twice as strong and tough as zirconia
- Zirconia is twice as strong and tough as alumina-based ceramics.

#### Which statement is correct: Fluorapatite is a fluoride containing: 10.

- а Calcium phosphate
- h Aluminium oxide
- Potassium oxide

# **CPD QUESTIONNAIRE**

## Article: Diagnosis 2013: the things you need to know for successful endodontic treatment. Thomas Jovicich, page 32

- 11. According to the author, the key component to endodontic treatment is diagnosis, using a multifocal approach that involves:
- Clinical signs and symptoms
- Diagnostic testing Radiographic findings None of the above All of the above d
- 12. According to the author, if the patient reports pain to bite upon release, this infers that there may be some structural root damage
- True
- False b
- 13. Which, according to the author, is the best way to test for cold symptoms:
- Refrigerant sprays
- Ice cubes (pellets) b
- CO2 systems

## 14. Which statement is correct:

- Pulp testing gold restorations can have the opposite effect to pulp testing porcelain restorations
- Pulp testing gold restorations can have the same effect as pulp testing porcelain restorations
- 15. According to the author, the best barometer to assess periodontal health, is probing from buccal to lingual with at least:
- Two measurements per side
- Three measurements per side b
- Four measurements per side

## Article: Interdisciplinary approach in aesthetic dentistry. Sebastian Ercus, page 44

- 16. According to the author, the correct ratio between the width and length of a maxillary central incisor is:
- 38 to 40 per cent
- 78 to 80 per cent b
- 58 to 60 per cent C.
- 17. The average distance from the lower margin of the upper lip and the base of the nose (subnasion) is.
- 11mm а
- 28mm h
- 21mm C

## 18. Which is incorrect. The '42.2 rule' refers to:

- A maximum of 4 mm of maxillary central incisor display when the lips are at rest а
- A minimum of 2 mm of maxillary central incisor display when the lips are at rest h
- A maximum of 4 mm of gingival display during smiling
- 19. In the case described, the aetiology of the excessive gingival display was a combination of :
- Delayed eruption
- h Altered passive eruption
- Hypermobility of the upper lip.
- None of the above
- e All of the above
- 20. In the case described, following the gingivectomy, a biologic width of:
- 4mm was maintained
- 2mm was maintained b
- 3mm was maintained C.