

CPD QUESTIONNAIRE 5.6.1

Article: Implant-prosthetic restorations – the challenge of creating an aesthetically pleasing smile in an edentulous patient. Petri, page 22

1. For patients desiring an alternative to complete dentures, which treatment option is becoming a frequent choice.
 - a Removable implant-supported overdentures
 - b Full-arch fixed bridge
2. Which statement is correct: In the case described, the overdenture was retained using:
 - a Bar attachments
 - b The double-crown method
3. How many implants were inserted to retain the overdenture prosthesis?
 - a Four
 - b Six
4. Which statement is correct. The primary telescope crowns were milled from:
 - a Lithium Disilicate
 - b Zirconia
 - c Gold
5. If the monomer concentration is reduced to less than 1% during the polymerisation process, the benefit to the patient is that the risk of allergies and irritation of the mucous membrane is virtually eliminated:
 - a True
 - b false

Article: How successful and predictable are implants placed with flapless guided surgery in fully edentulous rehabilitations? Hartshorne, page 30

6. Surgical and prosthetic complications during flapless guided implant surgery can be expected depending on the amount of deviation created by inaccuracies or error occurring from:
 - a Improper positioning of surgical guides during surgery
 - b 3D imaging
 - c Neither of the above
 - d Both of the above
7. Which statement is correct:
 - a Scott's pi statistic was used to measure the agreement of searches made by the two reviewers
 - b. Cohen's kappa statistic was used to measure the agreement of searches made by the two reviewers
8. The the most commonly reported complications in the study included:
 - a. High marginal bone loss
 - b. Prosthesis failure
9. Which statement is correct. During the scanning process, possible sources of deviation and error (inaccuracies) include:
 - a Incorrect positioning of the scan prosthesis
 - b Malpositioning and stabilization of the surgical template
 - c Patient movement during the scanning process
 - d None of the above
 - e All of the above
10. What, according to the author, is other catalyst that is important for the continued expansion and adoption of the digital workflow technology:
 - a CAD/CAM design software
 - b CBCT imaging technology
 - c The Dental Technician

CPD QUESTIONNAIRE 5.6.2

Article: Non-surgical endodontic retreatment of extensive periapical lesion. Ayón et al, page 54

11. Which of the following is not a cause of failure in endodontic treatment?
 - a Reinfection of the root canal by coronal leakage
 - b Foreign body reaction
 - c Failure to use calcium hydroxide as an interim dressing
 - d Extraradicular infection
12. Which is correct: The most widely used intracanal medication to date continues to be:
 - a Iodoform (CHI₃)
 - b Calcium hydroxide Ca(OH)₂
 - c Neither of the above
 - d Both of the above
13. In the case report described, which tooth was presented for treatment?
 - a The right lateral incisor
 - b The left lateral incisor
14. Which statement is correct:
 - a Lesions of 10mm or more are considered apical lesions
 - b The mean diameter of granulomas or apical cysts range from 5-8mm
 - c The mean diameter of apical lesions range from 5-8mm
15. Which statement is correct. Surgical procedures come with many drawbacks, which include:
 - a Pain and discomfort
 - b Possible damage to adjacent vital teeth and anatomic structures
 - c Neither of the above
 - d Both of the above

Article: Making sense of mouth ulceration: ulceration of local cause. Scully, page 62

16. Which of the following does not cause trauma:
 - a Assaults
 - b Self-harm (factitious).
 - c Bites
 - d Dental disease
17. A suspicious lesion should be observed for a period of:
 - a Two months
 - b Six weeks
 - c Two weeks
 - d Six months
18. Risk factors for radiation mucositis include:
 - a Concurrent chemotherapy
 - b Alcohol
 - c Poor oral hygiene
 - d None of the above
 - e All of the above
19. Traumatic ulcerative granuloma with stromal eosinophilia (TUGSE) is most commonly seen:
 - a On the lips
 - b On the buccal mucosa
 - c On the dorsal and lateral tongue
20. Which statement is correct.: Necrotising sialometaplasia (NSM) is:
 - a A rare condition of ulceration
 - b A common condition of ulceration