

interafrika  
dental  
pty(ltd)



biotène®  
oralbalance®

**THE COMPLETE DRY MOUTH TREATMENT**  
Boosts the mouth's natural defense system  
found in saliva



  
**Laclede, Inc.**  
HEALTHCARE PRODUCTS

Head Office: 1315 Church Street, Hatfield, Pretoria  
T: +27 12 342 8551/2/6  
F: +27 12 342 8541  
P O Box 11050, Hatfield, 0028, South Africa  
iad@mweb.co.za • www.iad.co.za

- 1 **Dental anxiety was found to be most prevalent in Australian women in the age group:**
  - a 26 - 35
  - b 35 - 44
  - c 44 - 60
  - d 60 +
- 2 **Which factors have been consistently linked with a greater incidence of dental anxiety?**
  - a Fear of pain
  - b Blood-injury fears
  - c Personality characteristics
  - d Past traumatic dental experiences
  - e All of the above
  - f None of the above.
- 3 **The study by Arntz revealed highly anxious patients:**
  - a experienced more pain than they expected
  - b experienced less pain than they expected
  - c expected and experienced no pain
- 4 **Studies have shown the most potent triggers of dental anxiety include:**
  - a Sight, sound and sensation of rotary dental drills
  - b Sight of dentist and staff
  - c Sight and sensation of dental local anaesthetic injection
  - d a and c
  - e None of the above
- 5 **Compared with non-anxious patients, dentally anxious patients have been found to have significantly:**
  - a Fewer missing teeth
  - b More filled teeth
  - c More missing teeth
  - d Fewer filled teeth
  - e a and b
  - f c and d
- 6 **The relationship between dental anxiety and satisfaction could result in the patient being:**
  - a Dissatisfied with the appearance of their teeth
  - b Less positive about their dentist
  - c None of the above
  - d Both of the above
- 7 **Upon identification of an anxious patient, the dentist should**
  - a Dismiss the patient's fears
  - b Ignore the patient
  - c Give the patient moral support
  - d Reassure the patient
  - e a and b
  - f c and d
- 8 **Complex distraction measures such as virtual reality techniques have been found to show:**
  - a more positive results for children than adults
  - b less positive results for children than adults
  - c the same results for children and adults
- 9 **Alternative methods for tooth preparation and caries removal include:**
  - a air abrasion
  - b ART (atraumatic restorative technique)
  - c chemo-mechanical caries removal (Carisolv®)
  - d all of the above
  - e none of the above
- 10 **Dental anxiety is a common problem found throughout the world, with more severe forms of fear and anxiety found in:**
  - a one in ten adults
  - b one in six adults
  - c one in a hundred adults

