

# Laser versus conventional therapies

Cristiane Meira Assunção,<sup>1</sup> Joanna Tatith Pereira,<sup>1</sup> Renata Schlesner Oliveira<sup>1</sup> and Jonas de Almeida Rodrigues<sup>2</sup>

## Introduction

In recent years, several studies have been conducted on the clinical applications of laser in dentistry. At the same time, there has been a marked emergence of organisations in support of the use of laser in dentistry. In the last decades, laser therapy has been used in dentistry as an adjunct or alternative to conventional approaches. In this paper, the following topics will be reviewed: the application of laser in caries prevention and diagnosis, hard- and soft-tissue treatments, and periodontal and endodontic procedures. There is a large research effort into new indications for laser in dentistry. It is expected that laser will become an essential component of the dentist's armamentarium.

While the technology was regarded as complex and of limited use in clinical dentistry in the past, a growing awareness of the usefulness of laser in the modern dental practice has been observed. Laser can be used as an adjunct or alternative to conventional approaches.<sup>1</sup> When comparing the use of laser with conventional therapies, three important areas must be considered: safety, efficacy and effectiveness. From an ethical standpoint, it is important to use the best available evidence when making clinical decisions.<sup>2</sup>

## Diagnostic laser applications

The most common methods for caries detection are visual and radiographic examination.<sup>3</sup> However, visual examination is a subjective method that depends on the knowledge and clinical experience of the examiner.<sup>3,6</sup> Several studies have demonstrated that radiographic examination demonstrates poor sensitivity to non-cavitated lesions.<sup>3,7,9</sup>

For this reason, fluorescence-based methods have been developed, aiming at the detection of occlusal and approximal carious lesions, for example DIAGNOdent 2095 (KaVo; LF; Figs. 1a-c) and DIAGNOdent 2190 (LF pen; Figs. 2a & b). They rely on the same principle: a laser diode emits red light at 655nm and a photodetector quantifies the reflected fluorescence from bacterial metabolites (fluorophores) in carious lesions, showing values ranging from 0 to 99.<sup>3,9</sup>

A study that assessed the performance of a visual method, radiographic examination and fluorescence-based methods in detecting occlusal caries in primary teeth found that the visual method and VistaProof fluorescence camera (Dürr Dental; FC) exhibited better accuracy in detecting enamel and dentine carious lesions, whereas the visual method combined with LF, LF pen and FC better detected dentine lesions on occlusal surfaces in primary teeth, with no statistically significant difference among them.<sup>3</sup>

Another study compared the performance of fluorescence-based methods (FC, LF and LF pen), radiographic examination, and another visual method called the International Caries Detection and Assessment System (ICDAS) II on occlusal surfaces. This study demonstrated that the combination of ICDAS and bite-wing radiographs yielded the best performance for detecting caries on occlusal surfaces.<sup>9</sup>

## Caries prevention: Enhancing enamel resistance

In the past, several in vitro studies have shown that enhancing enamel demineralisation resistance can be achieved by irradiation with lasers. In a blind in vitro study, Ana et

<sup>1</sup> Cristiane Meira Assunção,  
Joanna Tatith Pereira, Renata  
Schlesner Oliveira  
<sup>2</sup> Dr Jonas de Almeida  
Rodrigues, Brazil

Corresponding Author  
Dr Jonas de Almeida Rodrigues  
School of Dentistry, Federal  
University of Rio Grande do Sul  
Av. Paulo Gama, 110, Porto  
Alegre – RGS, 90000-000,  
Brazil  
jorodrigues@hotmail.com

al. 2012<sup>1</sup> compared the effect of professional fluoride application with that of laser irradiation with regard to the demineralisation of enamel and fluoride formation and retention. The study found that both methods enhanced enamel resistance, and no side-effects were found. A greater concentration of retained calcium fluoride-like material was found in the laser group. Formation and retention of calcium fluoride were also improved with laser irradiation.

The wavelengths absorbed most strongly by dental enamel are the 9.3 and 9.6µm carbon dioxide laser wavelengths. The reduction in acid dissolution of enamel is said to be caused by a loss of the carbonate phase of enamel crystals due to the heat of irradiation. Rechmann et al. 2011<sup>10</sup> demonstrated that short-pulsed 9.6µm carbon dioxide laser irradiation successfully inhibited enamel caries without any harm to the pulpal tissue of the teeth irradiated. The efficacy of carbon CO<sub>2</sub> laser irradiation regarding its long-term effect on caries resistances can be verified by further studies.

#### Hard-tissue applications: Caries removal

There is limited evidence to support the effectiveness of dental lasers in the removal of caries compared with rotary burs. In order to evaluate this, a systematic review of seven studies with adequate methodologies was performed.<sup>8</sup> Two of the studies found that there was no difference with regard to time taken for caries removal and cavity preparation. Four of the studies found that the laser took up to three times longer to perform these procedures. Four of the studies found that there were no differences between lasers and rotary burs with regard to pulpal effects. One of the studies found that dentists preferred the bur to the laser, and all the studies found that patients favoured the laser with respect to comfort. The studies found that adult patients prefer the laser, although the response from children was inconclusive. The results are not surprising, considering that local anaesthesia is often not needed when using a laser, making the overall dental experience more pleasant for the patient.<sup>10</sup>

#### Endodontic laser procedures (disinfection)

The main causes of endodontic treatment failure are the presence of persistent micro-organisms and recontamination of the root canal owing to inadequate sealing.<sup>11</sup> The long-term success rate of conventional endodontic treatment depends on several factors, such as the diverse and complex



Figure 1a: DIAGNOdent 2095.

anatomy of the root-canal system that consists of small canals diverging from the main canal. This complex system does not allow direct access during biomechanical preparation because of the canals' positioning and diameter.<sup>6</sup> New antimicrobial approaches to disinfecting root canals have been proposed; these include the use of high-power lasers and photodynamic therapy, which works by dose-dependent heat generation. However, in addition to killing bacteria, they have the potential to cause collateral damage such as charred dentine, ankylosed roots, melted cementum, root resorption and periradicular necrosis.<sup>2</sup>

In order to compare the effectiveness of antimicrobial photodynamic therapy with standard endodontic treatment and combined treatment to eliminate bacterial biofilms present in infected root canals, a study was conducted on ten single-rooted freshly extracted human teeth inoculated with stable bioluminescent Gram-negative bacteria. It found that endodontic therapy alone reduced bacterial bioluminescence by 90%, while photodynamic therapy alone reduced bioluminescence by 95%. The combination reduced bioluminescence by up to 98%, and, importantly, the bacterial regrowth observed 24 hours after treatment was much less for the combination group than for the treatment groups individually.<sup>12</sup>

Alternatives to conventional therapies to improve the disinfection of root canals are Nd:YAG and Er:YAG lasers. One study evaluated the bactericidal efficacy of Nd:YAG and Er:YAG lasers in experimentally infected curved root canals and concluded that in the straight root canals the



Figure 1b: Tip A for occlusal surfaces.



Figure 1c: Tip B for smooth surfaces.

Er:YAG laser had a bactericidal effect of 6.4 to 10.8% higher than that of the Nd:YAG laser. Conversely, the bactericidal effect of the Er:YAG laser in the curved root canals was 1.5 to 3.1% higher than that of the Nd:YAG laser.<sup>13</sup> These results suggest that further development of the endodontic laser tip and techniques are required to ensure its success.

### Periodontal laser procedures (disinfection)

Conventional periodontal therapy procedures include mechanical scaling and root planing, which has some limitations, especially in reducing bacteria inside deep pockets. In order to overcome the limitations of conventional mechanical therapy, several adjunctive protocols have been developed. Among these, laser has been proposed for its bactericidal and detoxification effects and for its ability to reach sites that conventional mechanical instrumentation cannot.<sup>14</sup>

Different lasers could be used in periodontal therapy for calculus removal, periodontal pocket disinfection, photoactivated dye disinfection of pockets and de-epithelialisation to assist regeneration.<sup>15</sup>

Several studies have indicated that the diode laser, with a wavelength of between 655 and 980nm, can accelerate wound healing through the facilitation of collagen synthesis, promotion of angiogenesis, and augmentation of growth factor release. Furthermore, the diode laser has in vitro bactericidal and detoxification effects and can prevent ablation of the root surface, which theoretically reduces the risk of removal of normal root tissue.<sup>13</sup>

Sgolastra et al. 2012<sup>14</sup> did not observe significant differences for any investigated outcome (clinical attachment level, probing depth, and changes in the plaque and gingival indices) in their systematic review. These findings suggest that

the use of the diode laser as an adjunctive therapy to conventional non-surgical periodontal therapy did not provide additional clinical benefit. However, given that few studies were included in the analysis, the results should be interpreted with caution. Important issues that remain to be clarified include the influence of smoking on clinical outcomes, the effectiveness of the adjunctive use of the diode laser on microbiological outcomes, and the effect of adverse events. Future studies are required to assess the effectiveness of the adjunctive use of the diode laser, as well as the appropriate dosimetry and laser settings.

### Soft-tissue applications

There are numerous soft-tissue procedures that can be performed with laser. Two key advantages of this are reduced intra-operative bleeding and less post-operative pain compared with conventional techniques, such as electrosurgery. Certain procedures in patients with bleeding disorders are better suited to lasers with greater haemostatic capabilities.<sup>5</sup>

### Conclusion

Although the results of laser therapy are similar (in safety, efficacy and effectiveness) to those obtained with conventional methods, new techniques and devices have been developed. Laser could thus be an evidence-based and well-supported treatment option for the dentist in daily dental practice.

### References

1. 15-Walsh, L. J. (2003). The current status of laser applications in dentistry. *Aust Dent J*, 48(3), 146-155; quiz 198.
2. 3-Dederich, D. N., & Bushick, R. D. (2004). Lasers in dentistry: separating science from hype. *J Am Dent Assoc*, 135(2), 204-212; quiz 229.
3. 14-Souza, J. F., Boldieri, T., Diniz, M. B., Rodrigues, J. A., Lussi, A., & Cordeiro, R. C. (2012). Traditional and novel methods for occlusal



Figure 2a: Cylindrical tip for occlusal surfaces.



Figure 2b: Wedge-shaped tip for proximal surfaces.

caries detection: performance on primary teeth. *Lasers Med Sci*.

4. 4-Diniz, M. B., Rodrigues, J. A., Hug, I., Cordeiro Rde, C., & Lussi, A. (2009). Reproducibility and accuracy of the ICDAS-II for occlusal caries detection. *Community Dent Oral Epidemiol*, 37(5), 399-404.

5. 15-Ekstrand, K. R., Martignon, S., Ricketts, D. J., & Qvist, V. (2007). Detection and activity assessment of primary coronal caries lesions: a methodologic study. *Oper Dent*, 32(3), 225-235.

6. 7-Jablonski-Momeni, A., Stachniss, V., Ricketts, D. N., Heinzel-Gutenbrunner, M., & Pieper, K. (2008). Reproducibility and accuracy of the ICDASII for detection of occlusal caries in vitro. *Caries Res*, 42(2), 79-87.

7. 9-Neuhaus, K. W., Rodrigues, J. A., Hug, I., Stich, H., & Lussi, A. (2011). Performance of laser fluorescence devices, visual and radiographic examination for the detection of occlusal caries in primary molars. *Clin Oral Investig*, 15(5), 635-641.

8. 11-Rodrigues, J. A., Diniz, M. B., Josgrillberg, E. B., & Cordeiro, R. C. (2009). In vitro comparison of laser fluorescence performance with visual examination for detection of occlusal caries in permanent and primary molars. *Lasers Med Sci*, 24(4), 501-506.

9. 12-Rodrigues, J. A., Hug, I., Diniz, M. B., & Lussi, A. (2008). Performance of fluorescence methods, radiographic examination and ICDAS II on occlusal surfaces in vitro. *Caries Res*, 42(4), 297-304.

10. 8-Makhija, S. K., & Anabtawi, M. F. There is limited evidence to support the effectiveness of dental lasers in the removal of caries compared with rotary burs. *J Evid Based Dent Pract*, 12(1), 31-32.

11. 16-Wong, R. (2004). Conventional endodontic failure and retreatment. *Dent Clin North Am*, 48(1), 265-289.

12. 6-Garcez, A. S., Ribeiro, M. S., Tegos, G. P., Nunez, S. C., Jorge, A. O., & Hamblin, M. R. (2007). Antimicrobial photodynamic therapy combined with conventional endodontic treatment to eliminate root canal biofilm infection. *Lasers Surg Med*, 39(1), 59-66.

13. 17-Yasuda, Y., Kawamorita, T., Yamaguchi, H., & Saito, T. (2010). Bactericidal effect of Nd:YAG and Er:YAG lasers in experimentally infected curved root canals. *Photomed Laser Surg*, 28 Suppl 2, S75-78.



Figure 3: Infra-red laser therapy for treatment of a primary herpetic infection in an adolescent patient undergoing chemotherapy (Therapy XT, DMC).

14. 13-Sgolastra, F., Severino, M., Gatto, R., & Monaco, A. Effectiveness of diode laser as adjunctive therapy to scaling root planning in the treatment of chronic periodontitis: a meta-analysis. *Lasers Med Sci*.

15. -Ana, P. A., Tabchoury, C. P., Cury, J. A., & Zezell, D. M. Effect of Er,Cr:YSGG laser and professional fluoride application on enamel demineralization and on fluoride retention. *Caries Res*, 46(5), 441-451.

16. 2-Bahcall, J. K., Miserendino, L., Walia, H., & Belardi, D. W. (1993). Scanning electron microscopic comparison of canal preparation with Nd:YAG laser and hand instrumentation: a preliminary study. *Gen Dent*, 41(1), 45-47.

17. 10-Rechmann, P., Fried, D., Le, C. Q., Nelson, G., Rapozo-Hilo, M., Rechmann, B. M., et al. Caries inhibition in vital teeth using 9.6- $\mu\text{m}$  CO<sub>2</sub>-laser irradiation. *J Biomed Opt*, 16(7), 071405.

Reprinted with permission by Laser 01/13